

German Freshwater Gastrotricha

by A. Schmidt-Rhaesa (May 2006)

sources:

Schwank, P. (1990): Gastrotricha. Vol. 3/1 in Süßwasserfauna von Mitteleuropa, Schwoerbel, J. & Zwick, P. (eds.), Gustav Fischer Verlag, Stuttgart

CHAETONOTIDA PAUCITUBULATINA

CHAETONOTIDAE

Aspidiophorus paradoxus (Voigt, 1902)

Holstein, Berlin

Aspidiophorus polonicus Kisielewski, 1981

Eastern Hessen

Aspidiophorus oculifer Kisielewski, 1981

Eastern Hessen

Aspidiophorus schlitzensis Schwank, 1990

Eastern Hessen, Black Forest

Chaetonotus acanthocephalus Valkanov, 1937

Eastern Hessen

Chaetonotus aemilianus Balsamo, 1978

Eastern Hessen

Chaetonotus alatus Schwank, 1990

Eastern Hessen

Chaetonotus arquatus Voigt, 1903

Bavaria

Chaetonotus balsaminus Schwank, 1990

Eastern Hessen, Black Forest

Chaetonotus bisacer Greuter, 1917

Berlin

Chaetonotus brevispinosus Zelinka, 1889

Central Germany

Chaetonotus caricola Schwank, 1990

Eastern Hessen

Chaetonotus christianus Schwank, 1990

Eastern Hessen, Southern Germany (near Bodensee)

Chaetonotus chuni Voigt, 1901

Whole Germany

Chaetonotus cordiformis Greuter, 1917

Franken

Chaetonotus disiunctus Greuter, 1917

Berlin, Franken

Chaetonotus elegans Konsuloff, 1921

Eastern Hessen, Black Forest, Franken

Chaetonotus euhystrix Schwank, 1990

Eastern Hessen

Chaetonotus ferrarius Schwank, 1990

Eastern Hessen, Black Forest

Chaetonotus heideri Brehm, 1917

Bodensee, Black Forest, Franken, Eastern Hessen
Chaetonotus heteracanthus Remane, 1927
Bavaria
Chaetonotus hoanicus Schwank, 1990
Eastern Hessen
Chaetonotus hystrix Metschnikoff, 1865
Whole Germany
Chaetonotus illiesi Schwank, 1990
Eastern Hessen
Chaetonotus laroides Marcolongo, 1910
Hessen, Black Forest, Franken, Holstein
Chaetonotus larus (O.F. Müller, 1773)
Whole Germany
Chaetonotus linguaeformis Voigt, 1902
Northern Germany
Chaetonotus longispinosus Stockes, 1887
Holstein
Chaetonotus macrochaetus Zelinka, 1889
Whole Germany
Chaetonotus maximus (Ehrenberg, 1831)
Whole Germany
Chaetonotus mitraformis Greuter, 1917
Berlin
Chaetonotus multispinosus Grünspan, 1908
Württemberg, Franken
Chaetonotus murrayi Remane, 1929
Berlin
Chaetonotus mutinensis Balsamo, 1978
Eastern Hessen
Chaetonotus octonarius Stockes, 1887
Whole Germany
Chaetonotus oculatus Schwank, 1990
Eastern Hessen
Chaetonotus ophiogaster Remane, 1927
Berlin
Chaetonotus persetosus Zelinka, 1889
Whole Germany
Chaetonotus ploenensis Voigt, 1904
Holstein, Franken
Chaetonotus polyspinosus Greuter, 1917
Franken, Alps, Central Germany, Black Forest, Eastern Hessen
Chaetonotus poznaniensis Kisielewski, 1981
Eastern Hessen, Black Forest
Chaetonotus pratensis Schwank, 1990
Eastern Hessen
Chaetonotus pygmaeus Schwank, 1990
Northern Germany
Chaetonotus remanei Schwank, 1990
Berlin
Chaetonotus rotundus Greuter, 1917
Eastern Hessen

Chaetonotus schlitzensis Schwank, 1990
Eastern Hessen

Chaetonotus schultzei Metschnikoff, 1851
Central Germany, Alps

Chaetonotus similis Zelinka, 1889
Whole Germany

Chaetonotus simrothi Voigt, 1909
Eastern Hessen, Bodensee

Chaetonotus spinulosus Stockes, 1887
Whole Germany

Chaetonotus succinctus Voigt, 1902
Franken

Chaetonotus tenuis Grünspan, 1908
Franken

Chaetonotus tricuspидatus Schwank, 1990
Eastern Hessen

Chaetonotus uncinus Voigt, 1902
Holstein

Chaetonotus voigti Greuter, 1917
Baden, Eastern Hessen

Chaetonotus vorax Remane, 1927
Berlin

Heterolepidoderma gracile Remane, 1927
Holstein

Heterolepidoderma macrops Kisielewski, 1981
Eastern Hessen

Heterolepidoderma majus Remane, 1927
Holstein

Heterolepidoderma ocellatum (Metschnikoff, 1865)
Broadly distributed throughout Germany

Heterolepidoderma patella Schwank, 1990
Eastern Hessen

Heterolepidoderma pineisquamatum Balsamo, 1980
Eastern Hessen, Black Forest

Ichthydium bifasciale Schwank, 1990
Eastern Hessen

Ichthydium forcipatum (Voigt, 1901)
Holstein

Ichthydium forficula (Remane, 1927)
Holstein

Ichthydium monolobum (Brunson, 1950)
Eastern Hessen

Ichthydium podura (O.F. Müller, 1773)
Northern Germany, Alps

Ichthydium sulcatum (Stockes, 1887)
Eastern Hessen, Black Forest

Lepidochaetus zelinkai (Grünspan, 1908)
Whole Germany

Lepidodermella squamata (Dujardin, 1841)
Broadly distributed throughout Germany
Lepidodermella zelinkai (Konsuloff, 1914)
Eastern Hessen, Black Forest

Polymerurus nodicaudus Voigt, 1901
Holstein, Central Germany, Alps
Polymerurus rhomboides (Stokes, 1887)
Upper Rhein river, Franken
Polymerurus serraticaudus (Voigt, 1901)
Upper Rhein river, Holstein

Rhomballichthys punctatus (Greuter, 1917)
Franken

DASYDYTIDAE

Haltidytes festinans (Voigt, 1909)
Holstein
Haltidytes saltitans (Stokes, 1887)
Upper Rhein river

Dasydytes asymmetricus Schwank, 1990
Eastern Hessen
Dasydytes bisetosus (Thompson, 1891)
Upper Rhein river
Dasydytes chatticus (Schwank, 1990)
Eastern Hessen
Dasydytes dubius (Voigt, 1909)
Holstein
Dasydytes goniathrix (Gosse, 1851)
Holstein
Dasydytes ornatus (Voigt, 1909)
Holstein

Chitonodytes collini (Remane, 1927)
Berlin
Chitonodytes longisetosus (Metschnikoff, 1865)
Gießen, Upper Rhein river

Stylochaeta fusiformis (Spencer, 1890)
Alps
Stylochaeta longispinosa Greuter, 1917
Northern Germany
Stylochaeta scirtetica Brunson, 1950
Northern Germany, Eastern Hessen
Stylochaeta stylifera (Voigt, 1901)
Holstein

NEOGOSSEIDAE

Neogossea antennigera (Gosse, 1851)

Central Germany
Neogosseia voigti (Daday, 1904)
Holstein