


MEIOFAUNA MARINA

Biodiversity, morphology and ecology
of small benthic organisms

19



MEIOFAUNA MARINA

Biodiversity, morphology and ecology of small benthic organisms

Volume 19 · September 2011

pages 1–208, 117 figs., 10 tabs., 1 app.

Chief editors

- Andreas Schmidt-Rhaesa Zoological Museum, University Hamburg, Martin-Luther-King-Platz 3, D-20146 Hamburg, Germany
Phone: +49 (0)40 42838-3921, Fax: +49 (0)40 42838-3937
E-mail: andreas.schmidt-rhaesa@uni-hamburg.de
- Kai Horst George Deutsches Zentrum für Marine Biodiversitätsforschung, Senckenberg am Meer, Südstrand 44, D-26382 Wilhelmshaven, Germany
E-mail: kgeorge@senckenberg.de
- M. Antonio Todaro Dipartimento di Biologia Animale, Università di Modena e Reggio Emilia, Via Campi 213/d, I-41100 Modena, Italia
E-mail: todaro.antonio@unimore.it

Editorial board

- Werner Armonies Alfred-Wegener-Institut für Polar- und Meeresforschung, Wattenmeerstation List auf Sylt
Thomas Bartolomaeus Freie Universität Berlin, Germany
Susan Bell University of South Florida, Tampa, FL, U.S.A.
Marco Curini Galletti University of Sassari, Italy
Nicole Dubilier Max-Planck-Institut für Molekulare Mikrobiologie, Bremen, Germany
Peter Funch University of Århus, Denmark
Gerhard Haszprunar Zoologische Staatssammlung, München, Germany
Rony Huys Natural History Museum, London, England
Ulf Jondelius University of Uppsala, Sweden
Marianne K. Litvaitis University of New Hampshire, Durham, NH, U.S.A.
Reinhardt Møbjerg Kristensen Zoological Museum, University of Copenhagen, Denmark
Ken-Ichi Tajika Nihon University School of Medicine, Tokyo, Japan
Seth Tyler University of Maine, Orono, ME, U.S.A.
Magda Vincx University Gent, Belgium
Wilfried Westheide Universität Osnabrück, Germany

Meiofauna marina is published annually

Subscriptions should be addressed to the Publisher:

Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, D-81379 München, Germany

PERSONAL SUBSCRIPTION: 48.– Euro

INSTITUTIONAL SUBSCRIPTION: 96.– Euro

Fees for mailing will be added

Manuscripts should be addressed to the editors

Articles appearing in this journal are indexed and abstracted in

Biological Abstracts

BIOSIS Previews

Bibliographic information published by the Deutsche Nationalbibliothek

The Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie;

detailed bibliographic data are available in the Internet at

<http://dnb.d-nb.de>.

Copyright © 2011 by Verlag Dr. Friedrich Pfeil, München, Germany

All rights reserved.

Printed by Advantage Printpool, Gilching

ISSN 1611-7557

Printed in the European Union

Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, D-81379 München, Germany

Phone: +49 (0)89 742827-0 · Fax: +49 (0)89 7242772 · E-mail: info@pfeil-verlag.de · www.pfeil-verlag.de

The spermatozoon of *Thaumastoderma moebjergi* with the description of the sperm model for the family Thaumastodermatidae (Gastrotricha, Macrodasysida)

Roberto Marotta*, Maria Balsamo**, Marco Ferraguti*, Cristina Fondello*,
 Loretta Guidi** and M. Antonio Todaro***

Abstract

The spermatozoon of *Thaumastoderma moebjergi* (Gastrotricha, Macrodasysida, Thaumastodermatidae) was described, and its structural features were compared to those of the other thaumastodermatid sperm models so far described. Although many of the sperm characters were present in different macrodasysidan species, the following complete set of characters was only found in all the thaumastodermatid spermatozoa at now described: a tubular structure within the acrosome formed by a pile of disks, the mitochondria arranged inside the nuclear spring, the striated cylinder in the flagellum, and a perinuclear helix. Thus the family Thaumastodermatidae appears uniform as far as sperm characters are considered.

Key words: Ultrastructure, systematics, spermatozoa, Gastrotricha

Introduction

We have few data on fertilization modalities in Gastrotricha, but, as far as we know, it is internal, through copulation, or transmission of spermatophores or dermal impregnation (Ruppert 1991). It is known that internal fertilization is correlated to changes in sperm morphology aimed to improve the movement inside the body's liquids of the partner (Franzén 1956), so we might expect a more slender and filiform cell, often accompanied

by some kind of helical shape at least in some parts of the cell. If, however, in an evolutionary line, internal fertilization has been “invented” more than once, the modifications undergone by the sperm cells may be very different from one species to another, due to different evolutionary pressures. Among Gastrotricha it seems that this is the case (Schmidt-Rhaesa 2007).

Our knowledge of gastrotrich sperm models has considerably increased in the last years (Marotta et al. 2005). We now have data on sperm

* Dipartimento di Biologia, Università degli Studi di Milano, via Celoria 26, 20133, Milano, Italy;
 e-mail: marco.ferraguti@unimi.it

** Dipartimento di Scienze della terra, della Vita e dell' Ambiente (DiSTeVA), Università di Urbino 'Carlo Bo',
 Loc. Crocicchia, 61029, Urbino, Italy

*** Dipartimento di Biologia, Università di Modena e Reggio Emilia, via Campi 213/d, 41125, Modena, Italy

morphology of species belonging to 21 different genera from twelve families. At least one species for each of the eight macrodasyidan families has been investigated, and we have data on eight species from the only four chaetonotidan families with amphigonic reproduction. As the number of species investigated increased, it became more and more evident that it is not possible to define a general model for gastrotrich spermatozoa (compare Ferraguti & Balsamo 1995 with Balsamo et al. 1999 and Marotta et al. 2005). Among macrodasyidans, the spermatozoa, besides being all filiform and flagellated cells (with the exception of *Dactylopodola baltica*: Fischer 1996 and *Dolichodasys*: Ruppert & Shaw 1977) which however is a character common to many animal species with internal fertilization, do not show characters common to all the species investigated. For example, one of the most peculiar characters of macrodasyidan gastrotrich spermatozoa, i. e. the presence of one or more mitochondria surrounded by the nucleus is not present in the family Macrodasysidae and in *Xenodasys eknomios* (Guidi et al. 2009). Another macrodasyidan sperm character, the striated cylinder involving the flagellar axoneme is absent in the *Lepidodasys* species examined (Guidi et al. 2004), in one of the two *Urodasys* species (Guidi et al. 2007), in two *Crasiella* species (Guidi et al. 2010), in *Xenodasys eknomios* (Guidi et al. 2009) and in the three Turbanellidae studied (Balsamo et al. 2002), but is present in all the remaining twelve species examined.

Thaumastodermatidae is the most speciose family within Macrodasysida, with near 150 species (Hochberg 2001, Hummon & Todaro 2010, Hummon 2011) distributed in eight genera and two recognized subfamilies: *Acanthodasys* and *Diplodasys* (Diplodasyinae) vs. *Hemidasys*, *Oregodasys* (= *Platydasys*), *Pseudostomella*, *Ptychostomella*, *Tetranchyroderma* and *Thaumastoderma* (Thaumastodermatinae). It is worth to notice that recently *Hemidasys* has been considered extinct (Hummon & Todaro 2010). Phylogenetic relationships within the family are debated, with contrasting scenarios hypothesized on gross anatomy level data (e.g. Hochberg & Litvaitis


2000 vs. Hochberg & Litvaitis 2001 vs. Kieneke et al. 2008).

We have spermatological data on species of four out of the eight genera: *Acanthodasys aculeatus* (Guidi et al. 2003), *Diplodasys ankei*, *Pseudostomella etrusca*, *Tetranchyroderma* (three species) (Ferraguti & Balsamo 1995). It appears that the general structure of the spermatozoa in the family Thaumastodermatidae is pretty constant. In particular all sperm models have a more or less complex, helical acrosome containing a tubular structure made of piled hollow cylinders, a nuclear-mitochondrial region in which one or more mitochondria are surrounded by a helical nucleus, and a tail in which the flagellar axoneme is surrounded by a striated cylinder (sensu Ferraguti & Balsamo 1994). To assess to what extent this general model is valid for other members of the family, and in search for possible character ground pattern we have studied the spermatozoa of *Thaumastoderma moebjergi* Clausen, 2004. Within this framework the study bears supplementary relevance as a recent phylogenetic investigation based on molecular traits found *Thaumastoderma* to be one of the most basal taxon along the Thaumastodermatinae evolutionary line (Todaro et al. 2011).

Materials and methods

Marine sediment containing *Thaumastoderma moebjergi* Clausen, 2004 was collected in September 2007 during a two-week workshop held at the Sven Lovén Centre for Marine Sciences on Tjörnö, an island on the Swedish west coast (for details see Willems et al. 2009). Gastrotrichs were extracted from the sediment by the narcotization-decantation technique, using an isosmotic (7 %) magnesium chloride solution (Todaro & Hummon 2008). The fauna-containing supernatant was then poured directly into a 5-cm diameter Petri dish and scanned for specimens under a Wild M3 dissecting microscope set at 50× magnification. For optical microscopy, the gastrotrichs were removed with a micropipette from the Petri dish, fresh-

Fig. 1. Spermatozoa of *Thaumastoderma moebjergi*. **A.** Main portion of the acrosome with the coiled tubular structure surrounded by the helical ridge (arrow). Inset: the apical portion of the spermatozoon. **B.** Basal portion of the acrosome to show the connection between the acrosome and the nucleus and the tubular vesicle (*) surrounding the distal portion of the nucleus. **C–D.** The nucleo-mitochondrial region of the spermatozoon in longitudinal (**C**) and cross (**D**) section. Note the two mitochondria in sequence (arrow). **E.** Basal portion of the nucleus with the beginning of the flagellum. **F.** Proximal region of the flagellum in longitudinal section. Note the two layers



forming the striated cylinder: the external, denser, one (arrow) and the inner, larger and more transparent (arrowhead). **G.** Cross section of the flagellum at the distal region of the axoneme. Note the irregular aspect of the $9 \times 2 + 2$ structure. **H.** distal portion of the tail, where the striated cylinder appears “empty” and closed distally by a double “cork” (inset). Scale bars: A-F, 0.2 μm ; G-H and insets, 0.1 μm .

mounted on slides and observed using a Nikon Eclipse 80i equipped with Differential Interference Contrast (Nomarski). During observation, the animals were photographed with a DS-5M Nikon digital camera and measured with the Nikon NIS software. For the transmission electron microscopy, 10 specimens were fixed overnight in 2 % glutaraldehyde in a 0.1 M sodium cacodylate buffer (pH 7.4), and stored in a 0.1 M sucrose/sodium cacodylate buffer. Subsequently, the animals were washed for two hours in 0.1 M cacodylate buffer, postfixed for two hours in aqueous 0.1 % Osmium Tetroxide in 0.1 M cacodylate buffer, washed in distilled water, pre-stained in a saturated solution of uranyl acetate, dehydrated in a graded ethanol series, and embedded in Spurr's resin.

Thin sections were obtained with a Reichert Ultracut E microtome, stained with 2 % lead citrate, carbon-coated and observed with a JEOL 100SX electron microscope.

The three dimensional reconstructions of the sperm model have been performed using Rhinoceros® NURBS (Non-Uniform Rational B-splines) modelling for Windows starting from the TEM micrographs.

Results

The spermatozoon of *Thaumastoderma moebjergi* is a long (more than 50 μm), filiform cell formed, in sequence, by a complex acrosome, a nucleo-mitochondrial region (together 16 μm long) and a tail (Fig. 2A-F).

The acrosome is a long organule with a diameter decreasing from about 0.2 μm at the proximal end, close to the nucleus (Fig. 1B) to about 0.1 μm apically, where it terminates with a short (0.5 μm), dense, conical end (Fig. 1A, inset; 2C). The acrosome contains a coiled tubular structure formed by a pile of electrondense rings (Fig. 1A), containing, basally, a series of tightly packed vesicles (Fig. 1B). The coiled tubular structure is surrounded by a lateral expansion in the form of a ridge following the coils of the tube, with a pitch increasing from 0.6 μm basally to 1 μm apically (Fig. 1A, 2A). The ridge is not surrounded by an acrosomal membrane, but is formed by a dense granular material. The plasma membrane surrounding the acrosome is somewhat loose. The acrosome is basally hosted in a short concavity formed by the nucleus (Fig. 1B).

The nucleo-mitochondrial region is formed by the compact nuclear chromatin forming a sort of "spring", with a diameter increasing from about 0.3 μm apically to 0.5–0.6 μm basally (Fig. 1C-E, 2B). The inner space delimited by the nuclear spring is often apically empty, in the first one-two coils (Fig. 1B, 2D), but is basally filled by the mitochondria (Fig. 1E). Since this area is quite long, it is difficult to establish the exact number of mitochondria present: in one instance we have seen two mitochondria disposed in line, one after the other (Fig. 1C). The nuclear chromatin forming the spring is uniform apically (Fig. 1C), but clearly divided in two parts basally, one more, the other less, electrondense (Fig. 2E). The distal portion of the nucleus, the one devoid of inner mitochondria, is surrounded by a tubular "vesicle" (0.1 μm in diameter) containing a dense central core surrounded by an electron transparent space, following externally the coils of the nucleus for 2.5 gyres (perinuclear helix in Guidi et al. 2003) (Fig. 1B, 2A,D).

The nucleo-mitochondrial region is followed by the flagellum, containing a conventional $9 \times 2 + 2$ axoneme, surrounded by a striated cylinder (Fig. 1G, 2E-F). This last is formed by two layers close one to the other: an external one, thinner and more electrondense, and an internal one, thicker (25 nm) which is formed by a U-shaped structure coiled around the axoneme and forming with its major axis an angle of 80–90 degrees (Fig. 1F). The tight coiling of the U-shaped structure gives to the organule the appearance of a striated cylinder (Fig. 1F, bottom). The axoneme originates from a complex connecting piece, without any conventional basal body, and terminates distally in an irregular way, so as many sections with odd number of microtubules have been found (Fig. 1G). The axoneme ends distally well before the end of the striated cylinder, thus leaving part of it with an "empty" appearance (Fig. 1H, 2F). The striated cylinder terminates distally in a peculiar, double end piece, the extremity of the spermatozoon (Fig. 1H and inset; 2F).

Discussion

The spermatozoa of *Gastrotricha* belong to the modified type sensu Franzén (1956) and are extremely complex cells, with a considerable variation of their characters among the taxa, so as we

were able to attempt a phylogenetic analysis of the group based on sperm morphology only (Marotta et al. 2005). The spermatozoon of *Thaumastoderma moebjergi*, when compared to the other sperm models known in the phylum, appears to share some of the characters with species of different families: the mitochondria surrounded by the nuclear spring are present also in Lepidodasyidae (Guidi et al. 2004), Cephalodasyidae (Ferraguti & Balsamo 1994, Fischer 1994), Planodasyidae (Guidi et al. 2010) and Turbanellidae (Balsamo et al. 2002); the striated cylinder around the axoneme is present in Cephalodasyidae (Ferraguti & Balsamo 1994, Fischer 1994, Balsamo et al. 1999) and Macrodasyidae (Marotta et al. 2005). However, all the members of the family Thaumastodermatidae, thus also *T. moebjergi* share the following complete set of characters:

- a tubular structure within the acrosome, formed by a pile of disks
- the mitochondria arranged inside the nuclear spring
- the striated cylinder in the flagellum
- the perinuclear helix. This last character has been overlooked in our former investigation of *Pseudostomella etrusca* (Ferraguti & Balsamo 1995), but a more careful observation has allowed us to recognize this character also in this species.

Thus the family Thaumastodermatidae appears uniform as far as sperm characters are considered. Each of the characters reported above do varies in the different species of the family:

- The tubular acrosomal structure in *Acanthodasys aculeatus* is formed by electron-dense rings surrounding an electron-lucent axis (Guidi et al. 2003), whereas in other species of the family the tubular structure is formed by thinner disks each with the shape of a thin circular crown. In *Diplodasys ankei* (Balsamo et al. 1999) the tubular acrosomal structure is formed basally by electron-dense rings that apically become electron-dense disks. In the same species the tubular acrosomal structure ends laterally against the acrosomal membrane, whereas in *Pseudostomella etrusca* and in *Tetranchyroderma* sp. it ends basally.
- There is only one mitochondrion in *Diplodasys ankei* (Balsamo et al. 1999), in *Acanthodasys aculeatus* (Guidi et al. 2003), and in *Tetranchyroderma* sp. 1 (Ferraguti & Balsamo 1995), at least two in *Thaumastoderma moebjergi*, but

many in *Pseudostomella etrusca* (Ferraguti & Balsamo 1995).

- The striated cylinder can be monolayered and very thick as in *Tetranchyroderma* sp. (Ferraguti & Balsamo 1995), or thinner as in *Acanthodasys aculeatus*: Guidi et al. 2003), but is thinner and formed by two layers in *Thaumastoderma moebjergi*. The distal extremity of the striated cylinder may be empty (i.e. it continues behind the axoneme) for a long tract, as in *Acanthodasys aculeatus* (Guidi et al. 2003), *T. moebjergi* and one *Tetranchyroderma* species.
- The perinuclear helix can run for one gyre only around the nucleus, as in *Pseudostomella etrusca*, or for two and a half, as in *Thaumastoderma moebjergi*, or involve the whole nucleus, as in *Acanthodasys aculeatus* (Guidi et al. 2003) and *Diplodasys ankei* (Ferraguti & Balsamo 1995, Balsamo et al. 1999). Our unpublished material on various *Tetranchyroderma* species allowed us to identify the perinuclear helix, but not to determine its length. We were not able to ascertain the origin of the perinuclear helix. Balsamo et al. (1999) proposed that the helix is a basal extension of the acrosome, and we had some confirmations from scattered images of the complex gastrotrich spermiogenesis (as for *Acanthodasys aculeatus*, Guidi et al. 2003), but its thinness and its coiled arrangement in the Thaumastodermatidae did not allow us to understand its real nature.

To summarize, the sperm characters of the family Thaumastodermatidae are fairly constant: a tubular structure inside the acrosome, the position of mitochondria inside the nuclear spring, the striated cylinder surrounding the axoneme, and the perinuclear helix.

Some of the above mentioned characters are present also in other gastrotrich taxa, whereas the perinuclear helix, at the actual state of knowledge, is present only in the species of the family Thaumastodermatidae. Unfortunately we could not detect ultrastructural features to support the evolutionary scenario proposed recently on molecular ground, in which *Thaumastoderma* is hypothesized as basal to most Thaumastodermatinae. The homology of the perinuclear helix to the acrosome of the extremely aberrant spermatozoon of *Urodasys acanthostylis* (Macrodasyidae) (Balsamo et al. 2007) remains to be established.




Fig. 2. Spermatozoon of *Thaumastoderma moebjergi*. **A-B.** 3D reconstructions of the acrosome (**A**) and nucleus (**B**). The acrosome is shown in its more distal (top) and proximal (bottom) regions. Basally, the acrosome is connected to the nucleo-mitochondrial region. Note the tubular vesicle coiled around the nucleus. **C-F.** Schematic drawing showing the apical regions of the acrosome (**C**), the acrosome-nuclear connection (**D**), the base of the flagellum (**E**) and the extremity of the tail (**F**). Abbreviations: **a**, acrosome; **hr**, helical ridge; **tv**, tubular vesicle; **n**, nucleus; **m**, mitochondria; **sc**, striated cylinder; **mt**, microtubules; **fep**, terminal structure of the flagellum.

Acknowledgements

This research has been supported by a PRIN 2007 grant to MB, MF, LG, and RM. M. A. Todaro is thankful to the organizing committee of the meiofauna workshop at the Sven Lovén Centre for Marine Sciences at Tjärnö during which *Thaumastoderma moebjergi* was collected.

References

- Balsamo, M., E. Fregni & M. Ferraguti (1999). Gastrotricha. In: *Reproductive Biology of the Invertebrates*, vol. IX, part A, Adiyodi, K. G. & R. G. Adiyodi (eds.), pp. 171–192. J. Wiley, Chichester, UK.
- Balsamo, M., M. Ferraguti, L. Guidi, M. A. Todaro & P. Tongiorgi (2002). Reproductive system and spermatozoa of *Paraturbanella teissieri* (Gastrotricha, Macrotrichida): implications for the sperm transfer modality in Turbellariidae. *Zoomorphology* 121: 235–241.
- Balsamo, M., L. Guidi, L. Pierboni, R. Marotta, M. A. Todaro & M. Ferraguti (2007). Living without mitochondria: spermatozoa and spermatogenesis in two species of *Urodasys* (Gastrotricha, Macrotrichida) from dysoxic sediments. *Invertebr. Biol.* 126: 1–9.
- Ferraguti, M. & M. Balsamo (1994). Sperm morphology and anatomy of the genital organs in *Mesodasys laticaudatus* Remane, 1951 (Gastrotricha, Macrotrichida). *J. Submicrosc. Cytol. Pathol.* 26: 21–28.
- (1995). Comparative spermatology of Gastrotricha. In: *Advances in Spermatozoal Phylogeny and Taxonomy*, vol. 166, Jamieson B. G. M., J. Ausio & J. L. Justine (eds.). *Mém. Mus. Natn. Hist. Nat. Paris* 166: 105–117.
- Fischer, U. (1994). Ultrastructure of spermiogenesis and spermatozoa of *Cephalodasys maximus* (Gastrotricha, Macrotrichida). *Zoomorphology* 114: 213–225.
- Fischer, U. (1996). Ultrastructure of penetrated spermatozoa, ovary, and oogenesis of *Dactylopodola baltica* (Gastrotricha, Macrotrichida). *Invertebr. Reprod. Dev.* 29: 71–78.
- Franzén, Å. (1956). On spermiogenesis, morphology of the spermatozoon, and biology of fertilization among invertebrates. *Zool. Bijdr. Uppsala* 31: 355–482.
- Guidi, L., M. Ferraguti, L. Pierboni & M. Balsamo (2003). Spermiogenesis and spermatozoa in *Acanthodasys aculeatus* (Gastrotricha, Macrotrichida): an ultrastructural study. *Acta Zool.* 84: 77–85.
- Guidi, L., L. Pierboni, M. Ferraguti, M. A. Todaro & M. Balsamo (2004). Spermatology of the genus *Lepidodasys*: towards a revision of the family Lepidodasyidae (Gastrotricha, Macrotrichida). *Acta Zool.* 85: 211–221.
- Guidi, L., M. Ferraguti, M. A. Todaro, L. Pierboni & M. Balsamo (2009). Unusual spermatozoa and reproductive modalities of *Xenodasys ekmanos* (Gastrotricha: Xenodasyidae). *Ital. J. Zool.* 76: 165–172.
- Guidi, L., M. A. Todaro, M. Ferraguti & M. Balsamo (2010). Reproductive system of the genus *Crasiella* (Gastrotricha, Macrotrichida). *Helgol. Mar. Res.* 65: 175–185.
- Hochberg, R. & M. K. Litvaitis (2000). Phylogeny of Gastrotricha: a morphology-based framework of gastrotrich relationships. *Biol. Bull.* 198: 299–305.
- (2001). Macrotrichida (Gastrotricha): a cladistic analysis of morphology. *Invert. Biol.* 120: 124–135.
- Hummon, W. D. (2011). Marine Gastrotricha of the Near East: 1. Fourteen new species of Macrotrichida, along with a species described from Australia. *Zookeys* 94: 1–59.
- Hummon, W. D. & M. A. Todaro (2010). Analytic taxonomy and notes on marine, brackish-water and estuarine Gastrotricha. *Zootaxa* 2392: 1–32.
- Kieneke, A., O. Riemann & W. H. Ahlrichs (2008). Novel implications for the basal internal relationships of Gastrotricha revealed by an analysis of morphological characters. *Zool. Scr.* 37: 429–460.
- Marotta, R., L. Guidi, L. Pierboni, M. Ferraguti, M. A. Todaro & M. Balsamo (2005). Sperm ultrastructure of *Macrotrichida caudatus* (Gastrotricha: Macrotrichida) and a sperm-based phylogenetic analysis of Gastrotricha. *Meiofauna Marina* 14: 9–21.
- Ruppert, E. E. (1991). Gastrotricha. In: *Microscopic Anatomy of Invertebrates*, Vol. 4, Aschelminthes. Harrison, F. W. & E. E. Ruppert (eds.). Wiley-Liss, New York: 41–109.
- Ruppert, E. E. & K. Shaw (1977). The reproductive system of gastrotrichs. I. Introduction with morphological data for two new *Dolichodasys* species. *Zool. Scr.* 6: 185–195.
- Schmidt-Rhaesa, A. (2007). *The Evolution of Organ Systems*. Oxford University Press.
- Todaro, M. A. & W. D. Hummon (2008). An overview and a dichotomous key to genera of the phylum Gastrotricha. *Meiofauna Marina* 16: 3–20.
- Todaro, M. A., T. Kånneby, M. Dal Zotto & U. Jondelius (2011). Phylogeny of Thaumastodermatidae (Gastrotricha: Macrotrichida) Inferred from Nuclear and Mitochondrial Sequence Data. *PLoS ONE* 6(3): e17892. doi:10.1371/journal.pone.0017892
- Willems, W. R., M. Curini-Galletti, T. J. Ferrero, D. Fontaneto, I. Heiner, R. Huys, V. N. Ivanenko, R. M. Kristensen, T. Kånneby, M. O. MacNaughton, P. Martínez Arbizu, M. A. Todaro, W. Sterrer & U. Jondelius (2009). Meiofauna of the Koster-area, results from a workshop at the Sven Lovén Centre for Marine Sciences (Tjärnö, Sweden). *Meiofauna Marina* 17: 1–34.

MEIOFAUNA MARINA

Biodiversity, morphology and ecology of small benthic organisms

INSTRUCTIONS TO CONTRIBUTORS

Meiofauna Marina continues the journal **Microfauna Marina**. It invites papers on all aspects of permanent and temporary marine meiofauna, especially those dealing with their taxonomy, biogeography, ecology, morphology and ultrastructure. Manuscripts on the evolution of marine meiofauna are also welcome. Publication of larger reviews or special volumes are possible, but need to be requested for. *Meiofauna Marina* will be published once a year. All contributions undergo a thorough process of peer-review.

Manuscript format: Manuscripts must be in English with metric units throughout. All parts of the manuscript must be typed, double-spaced, with margins at least 2.5 cm. Number all pages. Submit original plus 2 copies to facilitate reviewing and editing. Online-submission of manuscripts via the *Meiofauna Marina* homepage (www.meiofauna-marina.com) will be possible soon, but one of the editors must additionally be notified by e-mail or mail.

Page 1: Cover page including title of the paper; name(s) and address(es) of author(s); number of figures and tables. Suggest up to 5 keywords not in the title, and a short running title of no more than 50 characters. Indicate to which author correspondence and proofs should be sent; include e-mail, phone and fax numbers for this person.

Page 2: Concise abstract summarizing the main findings, conclusions, and their significance.

Page 3 and following pages: The Introduction, usually a brief account of background and goals, must be titled. Subsequent sections also bear titles, usually Material and Methods, Results, Discussion, Acknowledgements and References, but these may vary to suit the content. Subsections may be subtitled (don't number subtitles).

Figure legends, tables, and footnotes (in that order) should follow on extra pages following the References.

Citations and references: Complete data for all published works and theses cited, and only those cited, must be listed in References in alphabetical order; include papers accepted for publication (Cramer, in press), but not those merely submitted or in preparation. In the text, cite works in chronological order: (Smith & Ruppert 1988, Cook et al. 1992, Ax 1998a,b). Cite unpublished data and manuscripts from one of the authors (Smith, unpublished) or other individuals (E. E. Ruppert, pers. comm.) with no entry in References. Consult BIOSIS for journal-title abbreviations.

Examples of reference style:

Pesch, G. G., C. Müller & C. E. Pesch (1988). Chromosomes of the marine worm *Nephtys incisa* (Annelida: Polychaeta). *Ophelia* 28: 157-167.

Fish, A. B. & C. D. Cook (1992). Mussels and other edible Bivalves. Roe Publ., New York.

Smith, X. Y. (1993). Hydroid development. In: *Development of Marine Invertebrates*, vol. 2, Jones, M. N. (ed.), pp. 123-199. Doe Press, New York.

Illustrations and data: In designing tables, figures, and multiple-figure plates, keep in mind the final page size and proportions: 140 mm wide and maximally 200 mm high. Figures may occupy one column (68 mm) or two columns (140 mm). Details of all figures (graphs, line drawings, halftones) must be large enough to remain clear after reduction; type should be 1.5 mm high after reduction. Please submit original line drawings; they will be reduced to final size by the publisher.

Copies (submitted as hard copies or online) must be sufficiently good for reviewers to judge their quality. Include a scale bar and its value in each figure (value may be stated in the legend); do not use magnification. Authors are encouraged to submit extra, unlabelled photographs or drawings (black and white or colour) to be considered for the back cover of the journal. For final publication, photographic prints must be mounted, leaving no space between multiple prints on a plate. Protect each figure with a tissue cover sheet, and keep all materials within the size of the manuscript sheets, for safe and easy mailing.

Digital images and charts must be of high quality and professionally built. For more information visit "www.pfeil-verlag.de/div/eimag.php". Even if photographs or line drawings are processed with graphics programs, original slides, negatives or drawings must always be submitted.

Scientific names: For all species studied, the complete scientific name with taxonomic author and date (e.g., *Hesionides arenaria* Friedrich, 1937) should be given either at the first mention in the text of the paper or in the Material and Methods, but not in the title or abstract. Thereafter, use the full binomial (*Hesionides arenaria*) at the first mention in each section of the paper, and then abbreviate (*H. arenaria*, not *Hesionides* unless referring to the genus). Names for higher taxa should refer to monophyletic units, not to paraphyla (use, e.g., Macrostomida or Dinophilidae but not designations such as Turbellaria or Archiannelida). International nomenclature conventions must be observed, especially the International Code of Zoological Nomenclature (IRZN). The Latin name of any taxon is treated as a singular noun, not a plural or an adjective. Strictly, a taxon should not be confused with its members (the taxon Cnidaria does not bear nematocysts, but cnidarians do). Avoid terms of Linnean classification above the genus level.

Submitting a diskette: To facilitate speed and accuracy of publication, authors should supply a diskette after acceptance of the manuscript. Authors should retain a computer file that corresponds exactly to the hard-copy manuscript. Use a single standard font, a single space between sentences, and a single tab to indent each paragraph; avoid justifying, hyphenating, etc. Specialized word-processing commands (except boldface, italics, superscript, subscript) will have to be stripped from the final file. Use italics for species and genus names only. Complete instructions for diskettes will be sent with notification of acceptance.

Proofs, reprints, charges: 20 reprints are free of charge. Color plates must be paid by the authors. Additional reprints can be ordered by the authors.

MEIOFAUNA MARINA

Biodiversity, morphology and ecology
of small benthic organisms

Volume 19

CONTENTS

Schmidt-Rhaesa, Andreas, M. Antonio Todaro and Kai H. George: Editorial	1
Boaden, Pat J. S.: Swedmark, Kristineberg and the Advancement of Meiobenthology: a 1960's Nexus.....	3
Polte, Sabrina and Andreas Schmidt-Rhaesa: Immunohistochemical investigations of the interstitial cnidarian <i>Halammohydra octopodides</i> (Hydrozoa).....	17
Ax, Peter: New marine interstitial Plathelminthes from the Bay of Biscay, France.....	33
Delogu, Valentina and Marco Curini Galletti: <i>Sabussowia ronaldi</i> sp. nov. (Platyhelminthes: Tricladida: Maricola), a new Mediterranean species and its life cycle	41
Curini-Galletti, Marco, Marco Casu and Tiziana Lai: On the <i>Pseudomonocelis agilis</i> (Schultze, 1851) complex (Platyhelminthes: Proseriata), with description of two new species	49
Hummon, William D. and Jennifer L. Kelly: <i>Turbanella amphiatlantica</i> , a new species of Gastrotricha (Macrodasyida) from eastern North America and northwest Europe.....	61
Rothe, Birgen Holger, Alexander Kieneke and Andreas Schmidt-Rhaesa: The nervous system of <i>Xenotrichula intermedia</i> and <i>X. velox</i> (Gastrotricha: Paucitubulatina) by means of immunohistochemistry (IHC) and TEM	71
Marotta, Roberto, Maria Balsamo, Marco Ferraguti, Cristina Fondello, Loretta Guidi and M. Antonio Todaro: The spermatozoon of <i>Thaumastoderma moebjergi</i> with the description of the sperm model for the family Thaumastodermatidae (Gastrotricha, Macrodasyida)	89
Bartsch, Ilse: Marine mites (Acari: Halacaridae) from deep-sea gas seeps in the Gulf of Mexico	97
Koller, Stefan and Kai Horst George: Description of a new species of <i>Zosime</i> Boeck, 1872 (Copepoda: Harpacticoida: Zosimeidae) from the Great Meteor Seamount, representing one of the few eurybathic Harpacticoida among the distinct plateau and deep-sea assemblages	109
George, Kai Horst: Revision of the taxon <i>Fultonia</i> T. Scott (Copepoda: Harpacticoida: Argestidae), including the (re)description of some species, discontinuation of the genus <i>Parargestes</i> Lang, and erection of Argestinae Por, subfam. nov.	127
Addis, Alberto, Antonello Floris and Marcella Carcupino: A new species of the genus <i>Isocletopsyllus</i> (Harpacticoida, Cletopsyllidae)	161
Bruch, Katharina, Thomas Glatzel and Gritta Veit-Köhler: <i>Isthmiocaris laurae</i> sp. nov. (Crustacea, Copepoda, Harpacticoida) from the Angola Basin – first deep-sea species of the genus with remarks on its copepodid development	173
Schulze, Corinna and Andreas Schmidt-Rhaesa: Organisation of the musculature of <i>Batillipes pennaki</i> (Arthrotardigrada, Tardigrada)	195