


MEIOFAUNA MARINA

Biodiversity, morphology and ecology
of small benthic organisms

20



MEIOFAUNA MARINA

Biodiversity, morphology and ecology of small benthic organisms

Volume 20 · February 2013

pages 1-96, 28 figs., 10 tabs.

Managing editor

Kai Horst George Deutsches Zentrum für Marine Biodiversitätsforschung, Senckenberg am Meer, Südstrand 44, D-26382 Wilhelmshaven, Germany
E-mail: kgeorge@senckenberg.de

Co-editor

M. Antonio Todaro Dipartimento di Biologia Animale, Università di Modena e Reggio Emilia, Via Campi 213/d, I-41100 Modena, Italia
E-mail: todaro.antonio@unimore.it

Editorial board

Werner Armonies Alfred-Wegener-Institut für Polar- und Meeresforschung, Wattenmeerstation List auf Sylt
Thomas Bartolomaeus Freie Universität Berlin, Germany
Susan Bell University of South Florida, Tampa, FL, U.S.A.
Marco Curini Galletti University of Sassari, Italy
Nicole Dubilier Max-Planck-Institut für Molekulare Mikrobiologie, Bremen, Germany
Peter Funch University of Aarhus, Denmark
Gerhard Haszprunar Zoologische Staatssammlung, München, Germany
Rony Huys Natural History Museum, London, England
Ulf Jondelius University of Uppsala, Sweden
Marianne K. Litvaitis University of New Hampshire, Durham, NH, U.S.A.
Reinhardt Møbjerg Kristensen Zoological Museum, University of Copenhagen, Denmark
Ken-Ichi Tajika Nihon University School of Medicine, Tokyo, Japan
Seth Tyler University of Maine, Orono, ME, U.S.A.
Magda Vincx University Gent, Belgium
Wilfried Westheide Universität Osnabrück, Germany

Meiofauna marina is published annually

Subscriptions should be addressed to the Publisher:

Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, D-81379 München, Germany

PERSONAL SUBSCRIPTION: 48.– Euro

INSTITUTIONAL SUBSCRIPTION: 96.– Euro

Fees for mailing will be added

Manuscripts should be addressed to the editors

Articles appearing in this journal are indexed and abstracted in

Biological Abstracts

BIOSIS Previews

Bibliographic information published by the Deutsche Nationalbibliothek

The Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie;
detailed bibliographic data are available in the Internet at
<http://dnb.dnb.de>.

Copyright © 2013 by Verlag Dr. Friedrich Pfeil, München, Germany

All rights reserved.

Printed by Advantage Printpool, Wefßling

ISSN 1611-7557

Printed in the European Union

Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, 81379 München, Germany

Phone: +49 (0)89 742827-0 · Fax: +49 (0)89 7242772 · E-mail: info@pfeil-verlag.de · www.pfeil-verlag.de

A new eye-bearing *Macrodasys* (Gastrotricha: Macrodasyyida) from Jamaica

M. Antonio Todaro* and Francesca Leasi**

Abstract

A new macrodasyyidan gastrotrich is described from fine-medium sand collected at Doctor's cave beach of Montego Bay, Jamaica. *Macrodasys ommatus* n. sp. is the first gastrotrich to be reported from Jamaica and the second described species in the genus to bear eye-spots. The shape of the frontal organ distinguishes the Jamaican species from its sibling *M. nigrocellus*: elongate and undulate without an accessory lateral chamber in the former vs. rather short with an accessory lateral chamber in the latter. The following combination of characters further distinguishes the new species from its congeners: up to 7 anterior adhesive tubes per side arranged in a transversal row, three pairs of lateral adhesive tubes equally spaced along the intestinal region, up to 21 ventro-lateral adhesive tubes per side, two of which arise along the posterior region of the pharynx.

Key words: meiofauna, biodiversity, benthos, taxonomy, Caribbean Sea

Introduction

The study is part of a larger research programme aimed at shedding light on the diversity and phylogeny of gastrotrich species of the Tropical North-Western Atlantic (TNWA), with a focus on Small Islands Developing States (SIDS), whose sedimentary habitats are under environmental pressure from rising sea levels/shoreline erosion. Several international groups of researchers, along with students, are surveying gastrotrichs on several islands in the South Floridian, Bahamian, Lesser Antilles & Central Caribbean ecoregions. Preliminary accounts of these and related research

can be found in Atherton & Hochberg (2012a,b), Hochberg & Atherton (2010, 2011) and Hummon (2010a). Teams including the first author have so far visited three islands: St. John in the US-Virgin islands, Jamaica, and Curaçao. Part of the information and/or material from the visited islands appear in several works (e.g., Hummon et al. 2010; Kånneby et al. 2012a, 2012b; Todaro et al. 2012). We describe here a new species in the family Macrodasyyidae encountered in 2011 during a survey on the northern shore of Jamaica; it is also the first gastrotrich to be reported from Jamaica and the and the eight species of *Macrodasys* described from the TNWA (cf. Hummon 2010).

* Department of Life Sciences, University of Modena and Reggio Emilia, via Campi 213/D, I-41125 Modena, Italy; Email: antonio.todaro@unimore.it

** Department of Biodiversity, Earth and Environmental Science, Academy of Natural Sciences of Drexel University, Philadelphia, PA 19103, USA

Material and methods

The sampling campaign took place in February 2011 and included 10 locations along the North and West coasts of Jamaica. The species described herein was found in sublittoral samples collected at 1–4 m water depth, at Doctor's cave beach, Montego Bay. About 1 liter of sediment was sampled by skin-diving, collected into 500 ml plastic jars and soon after brought to the field laboratory (Discovery Bay Marine Laboratory). Here, the specimens were extracted daily with the narcotisation-decantation technique using a 7 % magnesium chloride solution within one week from collection; the supernatant was poured into plastic Petri dishes (3 cm diameter) and scanned for gastrotrichs at 50× under a Wild M3 stereomicroscope (Todaro & Hummon, 2008). When located, each individual gastrotrich specimen was mounted on glass slides and observed *in vivo* with Nomarski differential interference contrast optics using a Zeiss Axio Scope.A1. During observation, the specimens were photographed with a DS-5M Nikon digital camera and measured using the Nikon NIS-F software. Two specimens were fixed in 95 % ethanol and stored for future DNA analysis. The description of the new species follows the standardized scheme by Hummon et al. (1993), whereas the locations of some morphological characteristics along the body are given in percentage units (U) of total body length measured from anterior to posterior. Granulometric analysis of the substrata was carried out according to Todaro et al. (2006). Mean grain size, sorting coefficient, kurtosis, and skewness were calculated by a computerised programme based on the equation of Seward-Thompson & Hails (1973).

Taxonomic account

- Order Macrodasysida Remane, 1925
[Rao & Clausen, 1970]
Family Macrodasysidae Remane, 1926
Genus *Macrodasys* Remane, 1924

Macrodasys ommatus new species (Figs. 1–3)

Diagnosis. A *Macrodasys* species with a relatively stout body, with trunk longer than the pharyngeal region; head bearing sensory pits; pharyngo-intestinal junction (PhIJ) at U40; anterior adhesive

tubes (TbA), ventral in one row of six to seven tubes on each side of body, near anterior margin of head; lateral adhesive tubes (TbL), three per side along the intestinal region; ventro-lateral adhesive tubes (TbVL), about 20–21 per side, one anterior and one at the pharyngeal-intestinal junction, the remainder along the intestinal region; posterior adhesive tubes (TbP), 14 per side at the base and along the tail; frontal organ elongate (77 µm long and 9 µm wide), roughly spindle-shaped with an undulating border, lacking a clear differentiation between a spermathecal and a seminal receptacle region; nozzle not clearly cuticularised; caudal organ large, roughly S-shaped with anterior portion of the glandulomuscular organ showing curved, broadly-tipped apex and about the same length of the posterior portion; copulatory tube straight.

Type material. The description of *Macrodasys ommatus* n. sp. is derived from 3 specimens, two adults and a single juvenile, all collected from the same location. The holotype, LT = 513 µm, is the adult shown in Figure 2, (International Code of Zoological Nomenclature, Article 73.1.1), after observation it was fixed in 95 % ethanol and kept in the meiofauna collection of MAT for future DNA analysis. Likewise stored in ethanol is the paratypic juvenile, LT = 253 µm, shown in Figure 3 (ICZN Article 72.4.5); the second observed adult is not longer extant.

Type locality. The sediment samples were collected on 17th February 2011 from Doctor's cave beach, Montego Bay, Jamaica (Lat. 18°29'12" N; Long. 77°55'46" W).

Etymology. The specific name alludes to the presence of eye-spots (Gr. *omma*, eye)

Ecology. Occasional in frequency of occurrence (10–30 % of samples), rare in abundance (fewer than 1 % of a sample); sublittoral at 1–4 m water depth in fine sand (2.271 phi), moderately sorted (0.743 phi); water temperature, 26 °C; salinity, 33 psu.

Description. Adult specimens up to 513 µm total body length; PhIJ at U40; body strap-shaped, widest at the anterior trunk, head bearing piston pits and a pair of brown-to-black eye spots; each pigmented spot is contained in a cup-like, grayish structure; posterior trunk tapering gradually

into a short tail; widths at head, mid-pharynx, trunk and base of tail as follows: 53, 62, 77, 21 μm at U05/U28/U55/U89, respectively; dorsal and lateral surface covered with sensory hairs more densely packed on head; epidermal glands not observed.

Ciliature. The ventral locomotory ciliature is in the form of a continuous field spanning from posterior of the TbA to the caudal tail. Several pairs of sensorial bristles (12–18 μm in length) are visible along the margins and dorsal side of the body; other cilia (8–15 μm in length) with a putative sensorial function emerge from around the head.

Adhesive tubes. TbA (7–10 μm long), in a ventral, single row, forming an arc of 6–7 tubes, on each side of the body, adjacent to the mouth; TbL (18–21 μm long), 3 per side along the intestinal region at U46, U58 and U74 respectively; TbVL (14–16 μm long), 20–21 per side, one anterior and one at the pharyngeal-intestinal junction (at U28 and U40 respectively), the remaining tubes emerge along the intestinal region more or less regularly spaced the one from the other; TbP (15–10 μm long), 14 per side at the base and along the tail. There is no distinct cut off between TbVL and TbP but a rather continuous switch between the two groups at around U90.

Digestive tract. Mouth terminal (16 μm in diameter), leading to a small buccal cavity (16 $\mu\text{m} \times 20 \mu\text{m}$) which opens into a 180–189 μm long and 32–34 μm wide pharynx; pharyngeal pores at U28; intestine increases in width from the pharyngeal-intestine junction to U57, and gradually narrows up to the posterior body end; anus ventral at U86.

Reproductive tract. Hermaphroditic; elongate testes with the anterior-most portion at U40 and with vasa deferentia opening ventrally, approximately at U67. Frontal organ elongate, roughly spindle shaped, with an undulating appearance, about 77 μm long and 9 μm wide, without a clear differentiation between the posterior spermathecal and the anterior seminal receptacle regions; nozzle not clearly cuticularised. Caudal organ large (about 113 μm in length), roughly S-shaped with anterior region of the glandulomuscular portion showing curved, broadly-tipped apex and about the same length of the posterior portion; copulatory tube straight. Ovary adjacent to spermatheca; eggs maturing and increasing in size anteriorly, the full-grown one (54 μm long \times 41 μm wide) centered at U53.




Fig. 1. *Macrodasys ommatus* n. sp., schematic drawing of an adult specimen. Habitus as seen from the ventral side; sensorial- and locomotor ciliature omitted. CO, caudal organ; T, tail; Eg, egg; FO, frontal organ; PhIJ, pharyngo-intestinal junction; PP, pharyngeal pores; PtP, Piston pit; TbA, anterior adhesive tubes; TbL, lateral adhesive tubes; TbVL, ventro-lateral adhesive tubes; Ts, testis. Scale bar = 100 μm

Remarks. The juvenile specimen reached 258.6 μm in total length with PhIJ at U41. It had piston pits and brown eye-spots along with 3 TbA and 6 TbVL per side; the tail was well developed and furnished with eight adhesive tubes along its




Fig. 2. *Macrodasyx ommatus* n. sp., adult specimen, DIC photomicrographs. **A.** anterior region showing the eyespots, buccal cavity and the pharynx. **B.** mid- and posterior trunk region, showing the testes (Ts), frontal organ (arrowhead) and caudal organ (arrow). **C.** mid- and posterior trunk region at different focal plane showing the adhesive tubes of the lateral and ventrolateral series. Scale bars = 50 μ m.




Fig. 3. *Macrodasys ommatus* n. sp., juvenile specimen, DIC photomicrographs. **A.** habitus, dorsal side, showing the insertion of the sensorial bristles. **B.** habitus, showing the internal structures. **C.** habitus, ventral side, showing the adhesive tubes. Scale bars = 50 μ m.

margins. Seven pairs of sensorial bristles were visible on the dorsal side.

Taxonomic affinities. *Macrodasys* is one of the most speciose genera within the order Macrodasysida; currently, it includes 36 species (Hummon 2010a,b, 2011; Hummon & Todaro 2010); among these, only *Macrodasys nigrocellus* Hummon, 2011 bears eye spots like *M. ommatus* n. sp. The new species from Jamaica shares with *M. nigrocellus* from the Red Sea additional characteristics such as

the number and arrangement of the TbA, the presence of 2 pairs of TbVL along the posterior region of the pharyngeal region and 3 pairs of TbL along the intestinal region. The general appearance and especially the shape of the tail contribute further to make the two species appear similar. However, the shape of the frontal organ, elongate and with an undulate edge without an accessory lateral chamber in *M. ommatus* n. sp. vs. rather short with an accessory lateral chamber in *M. nigrocellus* n. sp. clearly differentiate the two taxa.

To our knowledge, beside *M. nigrocellus* and *M. ommatus* n. sp. at least three additional populations of *Macrodasys* specimens with eyes have been recorded. One from the Maldives, *Macrodasys* sp. (see Gerlach 1961), one from Belize (R. Hochberg, personal communication) and one (or two?) from Bahamas (A. Kieneke, personal communication). Based on the available information, none of these animals can be identified as *M. nigrocellus* or to *M. ommatus* n. sp. In fact, the scanty description of the Maldivian specimens is insufficient to make possible comparison with any other eye-bearing *Macrodasys* while additional information on the specimens from Belize and the Bahamas are needed to definitely clarify their taxonomic status.

Acknowledgements

This work was mainly supported by a grant from the US-National Science Foundation (n. DEB 0918499) to Rick Hochberg; additional funding was met by the BIOTOME project (UNIMORE) Many thanks are due to the staff of the Discovery Bay Marine Laboratory for the assistance received during our stay in Jamiaca. We are indebted to Rick Hochberg and Alexander Kieneke for sharing with us their information on eye-bearing *Macrodasys* from the Caribbean Sea.

References

- Atherton, S. & R. Hochberg (2012a). *Acanthodasys paurocactus* sp. n., a new species of Thaumastodermatidae (Gastrotricha, Macrodasysida) with multiple scale types from Capron Shoal, Florida. *ZooKeys* 190: 81–94.
- Atherton, S. & R. Hochberg (2012b). *Tetranchyroderma bronchostylus* sp. nov., the first known gastrotrich (Gastrotricha) with a sclerotic canal in the caudal organ. *Marine Biology Research* 8: 885–892.
- Gerlach, S. A. (1961). Ueber Gastrotrichen aus dem Meeressand der Malediven (Indischer Ozean). *Zoologischer Anzeiger* 167: 471–475.
- Hochberg, R. & S. Atherton (2010). *Acanthodasys caribbeanensis* sp. n., a new species of Thaumastodermatidae (Gastrotricha, Macrodasysida) from Belize and Panama. *ZooKeys* 61: 1–10.
- Hochberg, R. & S. Atherton (2011). A new species of *Lepidodasys* (Gastrotricha, Macrodasysida) from Panama with a description of its peptidergic nervous system using CLSM, anti-FMRamide and anti-SCP. *Zoologischer Anzeiger* 250: 111–122.
- Hummon, W. D. (2010a). Marine Gastrotricha of the Caribbean Sea: a review and new descriptions. *Bulletin of Marine Science* 86: 661–708.
- (2010b). Marine Gastrotricha of San Juan Island, Washington, USA, with notes on some species from Oregon and California. *Meiofauna Marina* 18: 11–40.
- (2011). Marine Gastrotricha of the Near East: 1. Fourteen new species of Macrodasysida and a redescription of *Dactylopodola agadasys* Hochberg, 2003. *ZooKeys* 94: 1–59.
- Hummon, W. D. & M. A. Todaro (2010). Analytic taxonomy and notes on marine, brackish-water and estuarine Gastrotricha. *Zootaxa* 2392: 1–32.
- Hummon, W. D., M. A., Todaro, T. Kånneby & R. Hochberg (2010). Marine Gastrotricha of the Caribbean Sea. Proceeding of the XIV International Meiofauna Conference. Gent, Belgium 12–16 July (Abstract).
- Hummon, W. D., M. A. Todaro & P. Tongiorgi (1993). Italian marine Gastrotricha: II. One new genus and ten new species of Macrodasysida. *Bollettino di Zoologia* 60: 109–127.
- Kånneby, T., M. A. Todaro & U. Jondelius, (2012a). A phylogenetic approach to species delimitation in freshwater Gastrotricha from Sweden. *Hydrobiologia* 683: 185–202.
- Kånneby, T., M. A. Todaro & U. Jondelius (2012b). Phylogeny of Chaetonotidae and other Paucitubulatina (Gastrotricha: Chaetonotida) and the colonization of aquatic ecosystems. *Zoologica Scripta* doi:10.1111/j.1463-6409.2012.00558.x.
- Seward-Thompson, B. L. & J. R. Hails (1973). An appraisal of the computation of statistical parameters in grain size analysis. *Sedimentology* 20: 161–169.
- Todaro, M.A. & W. D. Hummon (2008). An overview and a dichotomous key to genera of the phylum Gastrotricha. *Meiofauna Marina* 16: 3–20.
- Todaro, M. A., F. Leasi, N. Bizzarri & P. Tongiorgi (2006). Meiofauna densities and gastrotrich community composition in a Mediterranean sea cave. *Marine Biology* 149: 1079–1091.
- Todaro, M. A., M. Dal Zotto, U. Jondelius, R. Hochberg, W. D. Hummon, T. Kånneby & C. E. F. Rocha (2012). Gastrotricha: A Marine Sister for a Freshwater Puzzle. *PLoS ONE* 7: e31740.

MEIOFAUNA MARINA

Biodiversity, morphology and ecology of small benthic organisms

INSTRUCTIONS TO CONTRIBUTORS

Meiofauna Marina continues the journal **Microfauna Marina**. It invites papers on all aspects of permanent and temporary marine meiofauna, especially those dealing with their taxonomy, biogeography, ecology, morphology and ultrastructure. Manuscripts on the evolution of marine meiofauna are also welcome. Publication of larger reviews or special volumes are possible, but need to be requested for. *Meiofauna Marina* will be published once a year. All contributions undergo a thorough process of peer-review.

Manuscript format: Manuscripts must be in English with metric units throughout. Use italics for species and genus names only.

The original manuscript should be sent to the Managing Editor, Kai Horst George (kgeorge@senckenberg.de).

Additional information is requested:

Page 1: Cover page including title of the paper; name(s) and address(es) of author(s); number of figures and tables. Suggest up to 5 keywords not in the title, and a short running title of no more than 50 characters. Indicate to which author correspondence and proofs should be sent; include e-mail, phone and fax numbers for this person.

Page 2: Concise abstract summarizing the main findings, conclusions, and their significance.

Page 3 and following pages: The Introduction, usually a brief account of background and goals, must be titled. Subsequent sections also bear titles, usually Material and Methods, Results, Discussion, Acknowledgements and References, but these may vary to suit the content. Subsections may be subtitled (don't number subtitles).

Figure legends, tables, and footnotes (in that order) should follow on extra pages following the References.

Citations and references: Complete data for all published works and theses cited, and only those cited, must be listed in References in alphabetical order; include papers accepted for publication (Cramer, in press), but not those merely submitted or in preparation. In the text, cite works in chronological order: (Smith & Ruppert 1988, Cook et al. 1992, Ax 1998a,b). Cite unpublished data and manuscripts from one of the authors (Smith, unpublished) or other individuals (E. E. Ruppert, pers. comm.) with no entry in References. Consult BIOSIS for journal-title abbreviations.

Examples of reference style:

Pesch, G. G., C. Müller & C. E. Pesch (1988). Chromosomes of the marine worm *Nephtys incisa* (Annelida: Polychaeta). *Ophelia* 28: 157-167.

Fish, A. B. & C. D. Cook (1992). Mussels and other edible Bivalves. Roe Publ., New York.

Smith, X.Y. (1993). Hydroid development. In: Development of Marine Invertebrates, vol. 2, Jones, M. N. (ed.), pp. 123-199. Doe Press, New York.

Illustrations and data: In designing tables, figures, and multiple-figure plates, keep in mind the final page size and proportions: 140 mm wide and maximally 200 mm high. Figures may occupy one column (68 mm) or two columns (140 mm). Details of all figures (graphs, line drawings, halftones) must be large enough to remain clear after reduction. Labels, scale bars etc. should preferably be added with a vector-graphic program (e.g., Illustrator) or in text layers (Photoshop). Please submit line drawings with high resolution (TIF, 1200 dpi or more); they will be reduced to final size by the publisher. Send the original vector files, if a vector-graphic program is used. For more information visit "www.pfeil-verlag.de/div/eimag.php".

Copies (submitted as hard copies or online) must be sufficiently good for reviewers to judge their quality. Include a scale bar and its value in each figure (value may be stated in the legend); do not use magnification. Authors are encouraged to submit extra, unlabelled photographs or drawings (black and white or colour) to be considered for the back cover of the journal.

Scientific names: For all species studied, the complete scientific name with taxonomic author and date (e.g., *Hesionides arenaria* Friedrich, 1937) should be given either at the first mention in the text of the paper or in the Material and Methods, but not in the title or abstract. Thereafter, use the full binomial (*Hesionides arenaria*) at the first mention in each section of the paper, and then abbreviate (*H. arenaria*, not *Hesionides* unless referring to the genus). Names for higher taxa should refer to monophyletic units, not to paraphyla (use, e.g., Macrostomida or Dinophilidae but not designations such as Turbellaria or Archiannelida). International nomenclature conventions must be observed, especially the International Code of Zoological Nomenclature (ICZN). The Latin name of any taxon is treated as a singular noun, not a plural or an adjective. Strictly, a taxon should not be confused with its members (the taxon Cnidaria does not bear nematocysts, but cnidarians do). Avoid terms of Linnean classification above the genus level.

Reprints, charges: The authors will receive a PDF file for personal use free of charge; high-resolution open-access PDF files for unlimited use may be ordered. Color plates must be paid by the authors. Additional reprints can be ordered by the authors.

MEIOFAUNA MARINA

Biodiversity, morphology and ecology
of small benthic organisms

Volume 20

CONTENTS

Kai Horst George: Faunistic research on metazoan meiofauna from seamounts – a review.....	1
Todaro, M. Antonio and Francesca Leasi: A new eye-bearing <i>Macrodasys</i> (Gastrotricha: Macrodasysida) from Jamaica.....	33
Kieneke, Alexander, Birgen H. Rothe and Andreas Schmidt-Rhaesa: Record and description of <i>Anandrodasys agadasys</i> (Gastrotricha: Redudasyidae) from Lee Stocking Island (Bahamas), with remarks on populations from different geographic areas.....	39
Kieneke, Alexander, Sarah Narkus, Rick Hochberg and Andreas Schmidt-Rhaesa: <i>Diplo-dasys rothei</i> n. sp. (Gastrotricha, Macrodasysida), a new marine gastrotrich species from the Bahamas.....	49
Priscila A. Grohmann: First record of <i>Protohydra</i> cf. <i>leuckarti</i> (Cnidaria, Hydrozoa) from South America, in the Campos Basin, Rio de Janeiro, Brazil, with comments on its taxonomy and a brief review of its bio-ecology and distribution.....	63
Alexei V. Tchesunov: Marine free-living nematodes of the subfamily Stilbonematinae (Nematoda, Desmodoridae): taxonomic review with descriptions of a few species from the Nha Trang Bay, Central Vietnam.....	71
Publication Review.....	95

